

Writhe and Twist Helicity of Quantum Vortex Systems

RENZO L. RICCA

Department of Mathematics & Applications, U. Milano-Bicocca, Italy
E-mail: renzo.ricca@unimib.it

ABSTRACT

Here we show how kinetic helicity, a fundamental invariant of topological fluid dynamics, can be equally defined for quantum vortices under the Gross-Pitaevskii equation and we discuss how the integral definition of helicity in terms of linking numbers admits interpretation in terms of writhe and twist contributions [1]. We use this decomposition to study the evolution of writhe and twist under direct numerical simulation of the GPE evolution of a Hopf link and the following cascade process produced by a series of reconnections [2]. Since for quantum systems helicity remains always zero, and under anti-parallel reconnection writhe remains conserved across each reconnection event [3], twist must always balance changes in writhe. We show how this naturally leads to a continuous reduction of writhe and twist (of opposite sign) towards small-scale unknotted, unlinked planar rings. Moreover we also show [4] how induction of twist may induce the production of new, secondary defects in the system. This may have useful applications in laboratory production of defect entanglement in condensates.

- [1] Moffatt, H.K. & Ricca, R.L. (1992) Helicity and the Călugăreanu invariant. *Proc. R. Soc. A* **439**, 411-429.
- [2] Zuccher, S. & Ricca, R.L. (2017) Relaxation of twist helicity in the cascade process of linked quantum vortices. *Phys. Rev. E* **95**, 053109.
- [3] Laing, C.E., Ricca, R.L. & Sumners, DeW.L. (2015) Conservation of writhe helicity under anti-parallel reconnection. *Nature Sci. Rep.* **5**, 9224.
- [4] Foresti, M. & Ricca, R.L. (2019) Defect production by twist induction as Aharonov-Bohm effect. *Submitted*.